

PUBLICATIONS LOCE 2000 – 2004

- Eves P, Layton C, Hedley S, Dawson RA, Wagner R, Morandini R, Ghanem G, Mac Neil S.
Characterisation of an in vitro model of human melanoma invasion based on reconstructed human skin.
Br J Dermatology 142, 210-222, 2000.
- Dupont F, Avalosse B, Karim A, Mine N, Bosseler M, Maron A, Van den Broek A, Ghanem GE, Burny A, Zeicher M.
Tumor-selective gene transduction and cell killing using an oncotropic autonomous parvovirus-based vector.
Gene Therapy 7(9), 790-796, 2000.
- Haycock JW, Rowe SJ, Cartledge S, Wyatt A, Ghanem G, Morandini R, Rennie IG, Mac Neil S.
 α -Melanocyte Stimulating Hormone reduces impact of proinflammatory cytokine and peroxide generated oxidative stress on keratinocyte and melanoma cell lines.
J Biol Chem 275, (21), 15629-36, 2000.
- Kinnaert E, Morandini R, Simon S, Hill H, Ghanem G, Van Houtte P.
The degree of pigmentation modulates the radiosensitivity of human melanoma cells.
Radiation Res 154(6), 497-502, 2000.
- Hedley SJ, Murray A, Siskay K, Ghanem G, Morandini R, Gawkrödger D, MacNeil S
 α -Melanocyte Stimulating Hormone (α -MSH) can reduce T-cell interaction with melanoma cells *in vitro*.
Melanoma Res 10, 323-330, 2000.
- Mac Neil S, Eves P, Richardson B, Molife R, Lorigan P, Wagner M, Layton C, Morandini R, Ghanem G.
Oestrogenic steroids and melanoma cell interaction with adjacent skin cells influence invasion of melanoma cells in vitro.
Pigment Cell Res 13 suppl.8, 68-72, 2000.
- Jiménez-Cervantes C, Olivares C, González P, Morandini R, Ghanem G, García-Borrón JC.
The Pro162 variant is a loss-of-function mutation of the human Melanocortin 1 Receptor gene.
J Invest Dermatol 117(1), 156-158, 2001.
- Morandini R, Boeynaems JM, Wérenne J, Ghanem G.
Tips and step-by-step protocol for the optimization of important factors affecting cellular enzyme-linked immunosorbent assay (CELISA).
J Immunoassay and Immunochemistry, 22(4), 299-321, 2001.
- Ghanem G, Loir B, Morandini R, Salès F, Liénard D, Eggermont A, Lejeune F.
For the EORTC Melanoma Group.
On the release and half-life of S100B protein in the peripheral blood of melanoma patients.
Int J Cancer 94, 586-590, 2001.
- Süli-Vargha H, Nádasi H, Mihala N, Morandini R, Ghanem G.
Synthesis and biological activity of double cyclic RGD containing peptides.
In: Peptides 2000, proceedings of the 26th European Peptide Symposium, sept 10-15, 2000 - Montpellier, France. J. Martinez and Fehrentz JA Edts- EDK éditions médicales et scientifiques Paris, France. pp 749-750, 2001.

Vachtenheim J, Novotna H, Ghanem G.

Transcriptional repression of the microphthalmia gene in melanoma cells correlates with the unresponsiveness of target genes to ectopic microphthalmia-associated transcription factor.
J Invest Dermatol 117, 1505-1511, 2001.

Stoitchkov K, Letellier S, Garnier JP, Bousquet B, Tzankov N, Morel P, Ghanem G,
le Bricomb T.

Melanoma progression and serum L-Dopa/L-Tyrosine ratio : a comparison with S100B.
Melanoma Res 12(3), 255-262, 2002.

Lugassy C, Kleinman HK, Fernandez PM, Patierno SR, Webber MM, Ghanem G,
Spatz A, Barnhill RL.

Human Melanoma Cell Migration Along Capillary-Like Structures In Vitro: A New
Dynamic Model for Studying Extravascular Migratory Metastasis.
J Invest Dermatol 119(3):703-704, 2002.

Zhu N, Eves PC, Katerinaki E, Szabo M, Morandini R, Ghanem G, Lorigan P, MacNeil S,
Haycock JW.

Melanoma cell attachment, invasion, and integrin expression is upregulated by tumor
necrosis factor α and suppressed by α -melanocyte stimulating hormone.
J Invest Dermatol 119(5), 1165-1171, 2002.

Sánchez Más J, Olivares Sánchez C, Ghanem G, Haycock J, Lozano Teruel JA, García-
Borrón JC, Jiménez-Cervantes C.

Loss-of-function variants of the human melanocortin-1 receptor gene in melanoma cells
define structural determinants of receptor function.
Eur J Biochem 269,1-9,2002.

Moustafa M, Szabo M, Ghanem G, Morandini R, Kemp EH, Mac Neil S, Haycock JW.

Inhibition of TNF- α stimulated NF- κ B/p65 in Human Keratinocytes by α -Melanocyte
Stimulating Hormone and Adenocorticotropic Hormone Peptides
J Invest Dermatol 119(6), 1244-1253, 2002.

Brouard MC, Saurat J-H, Ghanem G, Siegenthaler G.

Urinary excretion of epidermal-type fatty acid-binding protein and S100A7 protein in
patients with cutaneous melanoma.
Melanoma Res 12, 627-631, 2002.

Eves P, Haycock J, Layton C, Wagner M, Kemp H, Szabo M, Morandini R, Ghanem G,
Garcia-Borrón JC, Jimenez-Cervantes C, Mac Neil S.

Anti-inflammatory and anti-invasive effects of alpha-melanocyte-stimulating hormone in
human melanoma cells.
Br J Cancer 17, 89(10), 2004-2015, 2003.

Stoitchkov K, Letellier S, Garnier JP, Bousquet B, Tsankov N, Morel P, Ghanem G,
Le Bricon T.

Evaluation of the serum L-dopa/L-tyrosine ratio as a melanoma marker.
Melanoma Res 13(6), 587-593, 2003.

Lugassy C, Kleinman HK, Engbring JA, Welch DR, Harms JF, Rufner R, Ghanem G,
Patierno SR, Barnhill RL.

Pericyte-Like Location of GFP tagged melanoma cells: Ex vivo and in vivo studies of
Extravascular Migratory Metastasis.
Am J Pathol 164(4), 1191-1198, 2004.

Kinnaert E, Duez P, Morandini R, Dubois J, Van Houtte P, Ghanem G.

Cysteine but not GSH modulates the radiosensitivity of human melanoma cells by affecting both survival and DNA damage.
Pigm Cell Res 17, 1-6, 2004.